

# Microsurgical Resection of Brain Stem Ependymoma: 2-Dimensional Operative Video


Mirza Pojskić, MD\*, Vincent Nguyen, MD‡, Goran Lakičević, MD, PhD§, Kenan I. Arnautović, MD, PhD\*¶

\*Department of Neurosurgery, University of Marburg, Marburg, Germany;

‡Department of Neurosurgery, University of Tennessee Health Science Center,

Memphis, Tennessee; §Department of Neurosurgery, University Clinical Hospital

Mostar, Mostar, Bosnia and Herzegovina; ¶Semmes Murphey Neurologic and Spine Institute, Memphis, Tennessee



Watch now at <https://academic.oup.com/ons/article-lookup/doi/10.1093/ons/oz252>

**Correspondence:** Kenan Arnautović, MD, PhD, Semmes Murphey Neurologic and Spine Institute, 6325 Humphreys Blvd., Memphis, TN 38120. Email: [kenanarnaut@yahoo.com](mailto:kenanarnaut@yahoo.com)

The brainstem is a less-common location for ependymomas than the spinal cord where they are the most common adult intramedullary tumor.<sup>1-18</sup> In this first video case report in the peer-reviewed literature, we demonstrate microsurgical resection of a medulla oblongata ependymoma.

There are several case reports of medulla oblongata ependymomas<sup>1,3,5,6,13</sup> and a few series of spinal cord ependymomas that included cases of ependymomas of the cervicomedullary junction.<sup>9,10</sup> The goal of surgery was to stabilize the preoperative neurological function; favorable outcome is achieved in patients with good preoperative statuses and well-defined tumor boundaries.<sup>9</sup> Although gross total resection (GTR) provides the best overall outcome, it is most effective for classic grade

II tumors, but not grade I (myxopapillary) and ependymomas, which have a lower GTR rate.<sup>14,15</sup>

A 55-yr-old patient developed 4-extremity weakness and dysphagia. Pre-/postcontrast magnetic resonance imaging (MRI) revealed centrally located brainstem lesion situated at the lower half of the medulla oblongata. Surgery, performed by the senior author, was performed in the prone position with a small suboccipital craniectomy and C1 posterior arch removal, followed by pia opening and posterior midline myelotomy. Tumor was debulked, dissected from the white matter, and resected. Histology revealed ependymoma (World Health Organization grade II). Postoperative pre-/postcontrast MRI revealed total resection. The patient's neurological deficit completely resolved postoperatively.

Written consent was obtained from the patient.

**KEY WORDS:** Brain stem ependymoma, Medulla oblongata ependymoma, Brain stem tumor, Tumor, Ependymoma, Microsurgical resection

*Operative Neurosurgery* 0:1-2, 2019

DOI:10.1093/ons/oz252

Received, March 13, 2019. Accepted, May 31, 2019.

## Disclosures

The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

## REFERENCES

- Amatya VJ, Takeshima Y, Kaneko M, et al. Case of clear cell ependymoma of medulla oblongata: clinicopathological and immunohistochemical study with literature review. *Acta Neuropathol.* 2003;53(5):727-738.
- Cikla U, Baggott C, Baskaya M. Gross total resection of large cervical intramedullary ependymoma: demonstration of microsurgical techniques. *Turk Neurosurg.* 2014;24(5):763-764.
- DiLuna ML, Levy GH, Sood S, Duncan CC. Primary myxopapillary ependymoma of the medulla: case report. *Neurosurgery.* 2010;66(6):E1208-1209; discussion E1209.
- Giammattei L, Penet N, Parker F, Messerer M. Intramedullary ependymoma: microsurgical resection technique. *Neurochirurgie.* 2017;63(5):398-401.
- Hoshimaru M, Koyama T, Hashimoto N. [Microsurgery of cervical intramedullary ependymomas extending into the medulla oblongata]. *No Shinkei Geka.* 2000;28(6):517-522.
- Ito T, Ozaki Y, Nakagawara J, Nakamura H, Tanaka S, Nagashima K. A case of cervicomedullary junction tanycytic ependymoma associated with marked cyst formation. *Brain Tumor Pathol.* 2005;22(1):29-33.
- Klekamp J. Spinal ependymomas. Part 1: intramedullary ependymomas. *Neurosurg Focus.* 2015;39(2):E6.

8. Kucia EJ, Bambakidis NC, Chang SW, Spetzler RF. Surgical technique and outcomes in the treatment of spinal cord ependymomas, part 1: intramedullary ependymomas. *Neurosurgery*. 2011;68(1 Suppl Operative):57-63; discussion 63.
9. Li D, Hao SY, Wu Z, Jia GJ, Zhang LW, Zhang JT. Intramedullary medullocervical ependymoma—surgical treatment, functional recovery, and long-term outcome. *Neurol Med Chir (Tokyo)*. 2013;53(10):663-675.
10. Li TY, Chu JS, Xu YL, et al. Surgical strategies and outcomes of spinal ependymomas of different lengths: analysis of 210 patients: clinical article. *J Neurosurg Spine*. 2014;21(2):249-259.
11. Liu JK. Laser-assisted microsurgical resection of thoracic intramedullary spinal cord ependymoma. *Neurosurg Focus*. 2012;33(Suppl 1):1.
12. McCormick PC. Microsurgical resection of intramedullary spinal cord ependymoma. *Neurosurg Focus*. 2014;37(Suppl 2): Video 9.
13. Mizuno J, Nakagawa H, Inoue T, Kondo S, Hara K, Hashizume Y. Signet-ring cell ependymoma with intratumoral hemorrhage in the medulla oblongata. *J Clin Neurosci*. 2005;12(6):711-714.
14. Oh MC, Tarapore PE, Kim JM, et al. Spinal ependymomas: benefits of extent of resection for different histological grades. *J Clin Neurosci*. 2013;20(10):1390-1397.
15. Safaei M, Oh MC, Mummaneni PV, et al. Surgical outcomes in spinal cord ependymomas and the importance of extent of resection in children and young adults. *J Neurosurg Pediatr*. 2014;13(4):393-399.
16. Samii M, Klekamp J. Surgical results of 100 intramedullary tumors in relation to accompanying syringomyelia. *Neurosurgery*. 1994;35(5):865-873; discussion 873.
17. Sayegh ET, Aranda D, Kim JM, Oh T, Parsa AT, Oh MC. Prognosis by tumor location in adults with intracranial ependymomas. *J Clin Neurosci*. 2014;21(12):2096-2101.
18. Wani K, Armstrong TS, Vera-Bolanos E, et al. A prognostic gene expression signature in infratentorial ependymoma. *Acta Neuropathol*. 2012;123(5):727-738.

## Acknowledgments

The authors wish to thank Andrew J. Gienapp (Neuroscience Institute, Le Bonheur Children's Hospital and Department of Neurosurgery, University of Tennessee Health Science Center, Memphis, Tennessee) for technical and copyediting, preparation of the manuscript for publishing, and publication assistance.